

the creation of a more comprehensive epidemiological profile of rotavirus infection in Australia.

References

1. Kapikian AZ, Chanock RM. Rotaviruses. In: Fields BN, Knipe DM, Howley PM, editors. *Fields Virology*. 3 ed. Philadelphia: Lippincott-Raven, 1996:1657-1708.
2. Bishop RF. Natural history of human rotavirus infections. In: Kapikian AZ (ed). *Viral infections of the gastrointestinal tract*. 2nd ed. Marcel Dekker, New York, 1994:131-167.
3. Carlin JB, Chondros P, Masendycz P, Bugg H, Bishop RF, Barnes GL. Rotavirus infection and rates of hospitalisation for acute gastroenteritis in young children in Australia, 1993-1996. *Med J Aust*. 1998;169 (5): 252-6.
4. Masendycz PJ, Unicomb LE, Kirkwood CD, Bishop RF. Rotavirus serotypes causing severe acute diarrhoea in young children in six Australian cities, 1989 to 1992. *J Clin Microbiol*. 1994;32:2315-2317.
5. Palombo EA, Bishop RF. Genetic and antigenic characterisation of a serotype G6 human rotavirus isolated in Melbourne, Australia. *J Med Virol*. 1995;47:348-354.
6. Palombo EA, Clark R, Bishop RF. Characterisation of a "European-like" serotype G8 human rotavirus isolated in Australia. *J Med Virol* 1999;51 (in press).
7. Palombo EA, Bugg HC, Masendycz PJ, Coulson BS, Barnes GL, Bishop RF. Multiple-gene rotavirus reassortants responsible for an outbreak of gastroenteritis in central and northern Australia. *J Gen Virol*. 1996;77:1223-1227.

Enterovirus 71 outbreak in Western Australia associated with acute flaccid paralysis. Preliminary report.

Peter McMinn,¹ Ivan Stratov,¹ Gary Dowse²

There have been 6 cases of acute flaccid paralysis (AFP) identified in Perth, Western Australia since March 1999 (2 in March, 2 in April, 1 in May and 1 in June). All cases have been in children under 2 years of age. Four of these cases are associated with enterovirus shedding (throat and stool) and EV71 has been identified by neutralisation in 2 of the cases. Stool specimens from 4 of the cases have been submitted to the Polio Reference Laboratory in Melbourne. Poliovirus infection has been ruled out in the 2 cases with confirmed EV71 infection and is pending in 2 other AFP cases.

Three of the 6 cases have residual weakness 1-2 months after illness onset and 1 case remains in hospital with prolonged flaccid paralysis requiring ventilation.

In addition, 12 cases of aseptic meningitis associated with enterovirus shedding (1 CSF isolate, the other isolates from throat and stool) have been identified. These cases have occurred in association with a large epidemic of hand, foot and mouth disease (HFMD) in Perth and in rural areas of Western Australia. One of the AFP cases comes from Kalgoorlie.

An enterovirus has been isolated from skin lesions or throat swabs from 15 uncomplicated HFMD cases, and 3 of 4 skin isolates have been confirmed as EV71. The EV71 isolates come from individuals living across the Perth metropolitan area and Mandurah, suggesting that EV71 activity has been widespread in recent months.

The origin of this virus is unknown. Whilst the recent EV71 epidemics in South-East Asia suggest a possible source, the acute myelitis observed in the Perth cases is clearly different from the brainstem encephalitis which characterised the severely affected cases in those epidemics. Molecular epidemiological studies of the Perth EV71 isolates are underway.

Public health measures instituted at this stage have included widespread media coverage to raise public awareness, and information and fact sheets distributed to all general practitioners and acute hospitals. In addition, child-care centres across the State have been given information emphasising the importance of good hygiene, and that children with HFMD should be excluded. Paediatric hospitals in other States and Territories have been advised.

1. Corresponding author. Department of Microbiology, Princess Margaret Hospital for Children, Roberts Road, Subiaco WA 6008)
2. Communicable Disease Control Branch, Health Department of Western Australia